

RESEARCH

Can humans perceive the metabolic benefit provided by augmentative exoskeletons?

Roberto Leo Medrano^{1,3*}, Gray Cortright Thomas^{1,2,3} and Elliott J. Rouse^{1,3}

Abstract

Background: The purpose of augmentative exoskeletons is to help people exceed the limitations of their human bodies, but this cannot be realized unless people choose to use these exciting technologies. Although human walking efficiency has been highly optimized over generations, exoskeletons have been able to consistently improve this efficiency by 10% - 15%. However, despite these measurable improvements, exoskeletons today remain confined to the laboratory. To achieve widespread adoption, exoskeletons must not only exceed the efficiency of human walking, but also provide a perceivable benefit to their wearers.

Methods: In this study, we quantify the perceptual threshold of the metabolic efficiency benefit provided during exoskeleton-assisted locomotion. Ten participants wore bilateral ankle exoskeletons during continuous walking. The assistance provided by the exoskeletons was varied in two minute intervals while participants provided feedback on their metabolic rate. These data were aggregated and used to estimate the perceptual threshold.

Results: On average, participants were able to detect a change in their metabolic rate of 22.7% (\pm 17.0%) with 75% accuracy. This indicates that in the short term, wearers are not able to perceive the metabolic benefit from any modern augmentative exoskeletons.

Conclusions: If wearers cannot perceive the benefits provided by these technologies, it will negatively affect their impact, including long-term adoption and product viability. Future exoskeleton researchers and designers can use these methods and results to inform the development of exoskeletons that reach their potential.

Keywords: exoskeleton; metabolic rate; perception; psychophysics; biomechanics

Background

The purpose of augmentative exoskeletons is to help people exceed the limitations of their human bodies. These technologies apply mechanical assistance to the joints of the legs during locomotion, thereby reducing the physical demands on the wearer's neuromuscular system. The potential uses for these technologies are broad and impactful, including assisting people's abilities to walk, run, jump, and/or carry loads. Consequently, lower-limb exoskeletons may improve the mobility of people with disabilities, as well as those completing sustained, physically demanding activities (*e.g.* first responders, postal/supply chain workers, and military personnel, among others). Recently developed systems for human augmentation applications are untethered [1, 2, 3], lightweight [4, 5, 1], and powerful

[6, 7]. While recent work has been encouraging, an ongoing challenge has been quantifying the success of these systems; the quantification of an exoskeleton's ability to reduce the metabolic expenditure of walking (*i.e.* calories burned) has emerged as a focus of the field [8].

Exoskeleton researchers have focused on the reduction of metabolic rate because it is intuitive, measurable, and supported by previous research. State-of-the-art exoskeletons have consistently reduced the metabolic expenditure needed for walking by approximately 14% relative to not wearing an exoskeleton [9, 10, 2, 5, 11, 12, 13]. These exoskeletons apply powered assistance at either the ankle joint [9, 10, 5, 11, 12, 13] or hip joint [2] and implement control strategies that operate in tandem with the wearer to reduce their metabolic expenditure. Intuitively, if an exoskeleton is successful, the muscular effort required will be reduced, which should be reflected in an upstream reduction in the metabolic power required from the wearer.

*Correspondence: ejrouse@umich.edu

¹Department of Mechanical Engineering, University of Michigan, 48109 Ann Arbor, USA

³Robotics Institute, University of Michigan, 48109 Ann Arbor, USA
Full list of author information is available at the end of the article

In addition, metabolic expenditure can be objectively measured in a laboratory setting, meaning it does not have the challenge of quantification that plagues other potentially subjective metrics of success (*e.g.* comfort, stability, or preference, among others).

There is mounting evidence that humans may be able to ‘subconsciously’ perceive their metabolic rate, but it is not yet known whether these changes can be perceived *consciously*. Donelan *et al.* showed that people choose step widths that minimize their metabolic rate during walking [14]. Subsequently, Selinger *et al.* demonstrated that exoskeleton wearers can re-optimize their gait patterns to minimize their metabolic rate when manipulated externally (with an exoskeleton) [15]. That is, the resistance of a knee exoskeleton was varied to incur a metabolic penalty during normative walking patterns, and participants needed to modify their gait patterns to reduce the superimposed metabolic burden. Participants converged to non-normative gait patterns that minimized metabolic rate, but this optimization did not occur spontaneously outside the laboratory [16]. Since exoskeleton wearers choose gait patterns that reduce metabolic rate, we believe this indicates people have some ability to sense this quantity (or something correlated). However, since people do not spontaneously stay or return to their lowest rate, it suggests that exoskeleton wearers do not have conscious knowledge of their metabolic rate or its gradient.

Conscious perception is a critical part of decision making. For an exoskeleton to appear valuable to its potential wearer, it must provide an experience that illustrates this value. Furthermore, this value must offset the potential “costs” of exoskeleton use. For example, without an intuitive and perceivable understanding of value, potential users may be unlikely to adopt exoskeletons with known disadvantages (*e.g.* monetary cost, discomfort, or being unfashionable). Previous research in the field of management science has investigated the implications of perceived value in technology adoption; one relevant framework proposed by Davis is the Technology Acceptance Model (TAM) [17]. In this model, Davis found a significant correlation between the consciously perceived usefulness of software and users’ intent to adopt the software [17]. More recently, King and He found that this relation was generalizable across many different technologies [18], such as broadband internet [19], telemedicine [20], and smart watches [21]. Thus, when potential exoskeleton users, manufacturers, and others are weighing the choice to adopt or purchase an exoskeleton, the consciously perceived value must outweigh the price, weight, aesthetics, and other costs of wearing a lower-limb exoskeleton.

The field of psychophysics focuses on quantifying human perception broadly [22]; for example, sensing may involve perception of images, temperatures, sounds [23], or metabolic rate. Forced comparison between two stimuli (such as asking which of two lines appears longer) across different trials is a powerful method for determining how humans can perceive changes in stimuli. This pair of stimuli is composed of a reference, which usually remains constant across trials, and a comparison, which changes from trial to trial. By analyzing large numbers of these comparisons, a perceptual model can be built that encodes and quantifies people’s perceptual performance. The input to this model is the true difference between the reference and the comparison, and the output is a probability of the comparison being perceived as different from the reference. The model predicts that stimuli are accurately perceived when the difference between the stimuli is large, but that human perception becomes essentially random when the difference is small. These models are often visualized using a *psychophysical curve* [24], typically a sigmoid function. In general, a single psychophysical curve pertains to the specific reference stimuli about which the test is conducted.

The steepness of a psychophysical curve quantifies perceptual ability; namely, the smallest difference in stimuli that can be perceived reliably. Using a threshold for reliability of 75% [25], this delta is known as the *Just Noticeable Difference* (JND). The JND has been used to quantify meaningful differences in visual acuity [26], sound [27], taste [28], and weight [29]. Recently, wearable robotics researchers have begun to quantify the JND of various factors in the design and control of wearable robotic systems, including perception of prosthetic ankle stiffness by users [30] and clinicians [31], environment stiffness [32] and viscosity [33], electrical stimulation of the residual limb [34], and vibrations of an osseointegrated prosthesis [35].

In this study, we characterized exoskeleton users’ conscious perception of their metabolic rate during assisted walking by quantifying the JND of metabolic rate changes. Understanding the human perceptual ability to sense this change is important because it has emerged as the gold standard by which exoskeletons are designed, controlled, and assessed. If exoskeletons are developed to impact a metric that is not perceivable by the user, it will likely hinder widespread success. To this end, we indirectly imposed different metabolic rates sequentially during walking by adjusting the assistance provided from bilateral ankle exoskeletons. Simultaneously, we recorded whether users perceived their metabolic rate to have increased or decreased as the control strategy changed. We aggregated these data to estimate the JND for changes in

metabolic rate. The contribution of this work includes new fundamental knowledge of how metabolic rate can be sensed during locomotion and a new benchmark for future exoskeleton developers who desire perceivable impact on metabolic expenditure. In addition, these results underscore the need for new metrics of exoskeleton success that are aligned with the value and experience of the user.

Methods

Participants

In this study, ten able-bodied participants ($N = 10$, 2 female, 8 male; age = 22.5 ± 3.17 years; mass = 70.9 ± 11.9 kg, Table. 1) walked using bilateral ankle exoskeletons on a treadmill. The required number of participants was chosen based on a power analysis to quantify a JND of 15% with 80% power and 5% type 1 error rate. We chose 15% as this was representative of the reductions achieved by the best performing lower-limb exoskeletons [36, 37, 38]. All participants provided written informed consent before participation. The study protocol was approved and overseen by the Institutional Review Board of the University of Michigan Medical School.

Table 1 Participant Data

Participant	Number of Responses	Gender	Weight	Age
[0.5ex] 1	53	M	52.2	20
2	100	M	74.0	23
3	34	F	72.0	21
4	100	M	86.0	24
5	100	M	78.5	21
6	100	M	74.0	23
7	100	M	82.5	24
8	100	M	59.0	19
9	100	F	53.5	20
10	100	M	77.0	30

Experimental Protocol

Walking Protocol

Participants experienced numerous metabolic rate changes in sequence that stemmed from the assistance provided by the ankle exoskeletons. Participants walked for 20 minute blocks, where each block consisted of 10 trials in series. Following each pair of trials, participants responded regarding which condition they perceived had a higher metabolic rate by agreeing or disagreeing to the binary question “*is the current level of exertion higher than the previous level of exertion?*”. Participants responded non-verbally with either a ‘thumbs up’ or a ‘thumbs down.’ Thus, each block consisted of nine comparisons across ten trials and participants completed approximately 11 blocks across three to four days of data collection. We chose

the two minute walking duration for each trial to balance metabolic estimation quality with experiment duration; Zhang *et al.* demonstrated that the metabolic estimation error with two minutes of data is approximately 2% [39]. The two-minute trial duration also allowed the participants adequate time to experience and react to each walking condition.

Prior to the experiment, participants familiarized themselves with several aspects of the experimental protocol. Participants read a lay explanation of metabolic rate to familiarize themselves with the concept. Next, participants were primed to react to their feeling of general exertion by reading the instructions of the Borg Rating of Perceived Exertion [40, 41]. We chose to have the participants read information on the Borg Scale because it has previously been demonstrated to produce accurate estimates of exertion [42]. Finally, participants underwent a four-minute acclimatization period in which they were exposed to different representative exoskeleton behaviors that spanned what could be encountered during the experiment.




Figure 1 Sample exoskeleton current profiles used in this experiment (colored lines). The profiles resembled square pulses and were parametrized using the following parameters: the peak current of the profile, the onset time of the profile, and the duration of the profile. The total bounds of possible current profiles are delineated by the black dashed line. The profiles’ currents were mapped to motor torques through the motor torque constant and exoskeleton’s transmission.

Exoskeleton Control

We used bilateral ankle exoskeletons (Dephy ExoBoot, Dephy Inc. Maynard MA) to manipulate the metabolic rate of the wearer. The exoskeleton (Fig. 2B) used electric motors (~ 300 W) and flat cable transmission ($\sim 15:1$) to apply plantarflexion assistance during walking. The assistance was governed by parameterized

current profiles that resembled a square pulse (see Fig. 1). The current profiles were governed by three parameters; we manipulated the onset timing, pulse magnitude, and pulse duration. We chose i) onset timings from a uniform distribution bounded between 25% and 50% of stride time, ii) pulse magnitudes from a uniform distribution bounded between 15 A and 25 A (corresponding to approximately 12 and 20 Nm with the ExoBoot's nonlinear transmission), and iii) pulse durations from a uniform distribution with variable bounds. The variable bounds for the pulse duration depended on the sample drawn from the onset timing distribution such that the square pulse had a minimum duration of 10% of stride, and a maximum duration of 60% of stride time. Onset timings that occurred earlier than 30% of stride were additionally constrained to have a minimum pulse duration of 20% of stride, which was imposed to prevent excessive device wear. We chose these bounds as they have been shown to significantly alter participant metabolic rate, and thus allow us to sample as wide an energetic range as possible [43, 44, 12] while balancing device integrity and user safety. The current profiles were described using the stride completion percentage to mitigate any variations in step length or cadence that occurred during the trial. Thus, we inferred the stride completion percentage using heel-strike events. We detected these events by thresholding the onboard accelerometers (MPU-9250, InvenSense, San Jose, CA) [45, 46].

Metabolic Rate Sensing

Participants walked with a randomized torque profile for two minutes, which produced a first-order dynamic response in metabolic rate [47]. We measured participant metabolic rates through indirect calorimetry [48] (COSMED K5, Rome IT) (Fig. 2A). We estimated the user's steady state metabolic rate by fitting a first-order response [47, 43] to the breath-by-breath transient data gathered over the two minutes for each trial, with the steady state value representing the trial's metabolic rate. Prior to undergoing the walking protocol, participants stood still for four minutes to obtain their baseline metabolic rates. Each participant's standing metabolic rate was computed as the average rate over this four-minute interval. The standing metabolic rate was subtracted from each trial's metabolic rate measurement to isolate the metabolic effects of exoskeleton assisted locomotion (*i.e.* net metabolic rate).

Psychophysical Function Fitting

To estimate the Just Noticeable Difference (JND) of metabolic rate, which denotes the magnitude of change necessary for consistent perception, our experimental

protocol requires the normalization of metabolic rates. That is, to compare across sequential trials with differing references, normalization is needed to combine these data to obtain a single JND for each subject under the assumption of a constant Weber Fraction (see Limitations subsection) [49]. The Weber Fraction (WF) [50] is a metric that captures the differences in perceptual thresholds that are dependent on the magnitude of the reference stimulus used in the comparisons. By definition, the WF is the JND divided by the reference stimulus, thus it represents the percent change from the reference stimulus that is perceivable. For a wide range of stimulus magnitudes, the WF can be modeled by a constant [51].

For normalization, consider a sequential pair of metabolic rates A then B; we normalized B (the comparison) as a percent change in rate from A. Note that rate A is the reference that changes from trial to trial. We used the normalized metabolic rate differences and corresponding participant responses to fit a psychometric function. The psychometric function then provided JNDs with units of percent change of metabolic rate (rather than absolute units (W/kg)). The JND is then equivalent to the WF expressed as a percentage.

A logistic psychometric function was used to model participant responses. This model predicted the probability that the participant would choose "the comparison is *greater*" as a function of the normalized metabolic rate difference between the two trials. Using the convention from above, the psychophysical curve predicts the probability that rate B is greater than rate A, as a function of the relative difference between A and B. The logistic function of (reference-normalized) stimulus x had the following form,

$$\Psi(x, \alpha, \beta, \gamma, \lambda) = \gamma + \frac{1 - \lambda - \gamma}{1 + e^{-\beta(x - \alpha)}} \quad (1)$$

where $\Psi(x)$ was parametrized by the following variables: the experimental lapsing rate λ , which was fixed at the commonly used value of 0.02 [22]; the false positive rate γ , which was fixed at the lapsing rate since participants underwent a stimulus discrimination task [22, 52]; the logistic function's threshold point α on the x-axis, which anchors the center of the logistic curve and was set to 0; and the parameter β which governs the slope of the logistic function and is the only degree of freedom estimated during the fitting procedure.

Using the modeled psychometric curve, we quantified the Just Noticeable Difference (JND) which represents the minimum change in metabolic rate that must occur before an observer can reliably perceive with 75% accuracy [53]. The JND is calculated by taking the difference between the values of x at $\Psi(x) = 0.75$ and

$\Psi(x) = 0.25$ and dividing the difference by two. By fixing the other parameters of $\Psi(x)$ at the values specified, the JND thus depended only on β

$$\text{JND} = k/\beta, \quad (2)$$

with a scale constant k in terms of the fixed parameters,

$$k = \frac{1}{2} \ln \left[\frac{(0.75 - \gamma)(1 - \lambda - 0.25)}{(1 - \lambda - 0.75)(0.25 - \gamma)} \right]. \quad (3)$$

Shallower slopes (indicating less sensitivity) caused higher JNDs, while steeper slopes (indicating higher sensitivity) caused lower JNDs.

Statistics and Comparisons


A separate logistic model was fit for each participant using Bayesian analysis [54]. This approach yielded a posterior distribution of JND estimates for each participant. From this posterior distribution, we extracted the maximum likelihood estimate for each participant, which was considered the estimated JND [55]. Our approach of using Bayesian estimation enables quantification of both the JND value for each subject in addition to the uncertainty about our estimates. We chose Bayesian estimation because preliminary work indicated the conventional Maximum Likelihood Estimation approach could fail to converge [56]. We conducted our Bayesian analysis using the PyMC3 library in Python [57]. Each participant's prior distribution of JND estimates was chosen as a uniform distribution between 0% and 70%, representing a plausibly large range of perceptual abilities.

The posterior JND distributions were obtained by updating our prior distributions using the participant response data. We used the No-U-Turn Sampler (NUTS) [58] strategy—a Markov Chain Monte Carlo (MCMC) algorithm—to numerically approximate the posterior distribution of possible JND values; we used four sampling chains with 8000 tuning iterations and 4000 posterior predictive samples. We chose these values to balance computation time and accuracy. The JND with the highest likelihood in the posterior distribution was the nominal JND estimate. Each posterior also yielded a 95% credible interval for the JND estimates.


Our approach using Bayesian statistics enabled investigation of several assumptions made about the JND distributions. We compared three different JND models using the Watanabe-Akaike Information Criterion (WAIC) metric, which evaluates the predictive power of models and corrects for the number of model

parameters to favor parsimony [59]. The three competing models were: i) a pooled model that featured a single JND parameter and posterior distribution that applied to all participants; ii) an independent model that assumed each participant had a different JND; and iii) a variable Weber Fraction (WF) model that allowed for multiple JND estimates per participant depending on the magnitude of the reference data. In each model, our parameter estimates were informed by the data, yielding posterior distributions over all possible parameter estimates. In the pooled model, the single JND estimate predicted the responses of all participants; in the independent model, each participant's responses were predicted by individual JND distributions which we estimated; and in the the variable-WF model, each participant had two different JNDs for references above or below the average reference value. The pooled model featured 2000 tuning iterations and 2000 posterior predictive samples for reduced computational time, given the greater number of responses used as input. The remaining models used our default settings of 8000 tuning samples and 4000 posterior predictive samples due to the relative sparsity of the data and the complexity of the models.

Results



The average inter-participant JND was 22.7% (standard deviation (SD): 17.0%) (Fig. 3A). Many participants were highly attuned to the changes in their metabolic energetics, while others were less perceptive,



as evidenced by the high standard deviation of the estimates (Fig. 3B). We used a one-sample Kolmogorov-Smirnov test in MATLAB to verify that the independent JND estimates from the participants were normally distributed. The lowest estimated JND was 6.4%, while the highest was 69.9%. Our confidence in each participant's JND estimate was given by their respective JND posterior distributions, which represent the distribution over potential JNDs of each participant (Fig. 3C). Differences in the JNDs can then be observed by comparing the shapes of these distributions; for example, a narrow distribution with a defined peak at a low value represents a participant who is highly attuned to changes in energetics, while a flattened distribution with a peak at a high value denotes a participant with a greater JND and less sensitivity.

We used the Watanabe-Akaike Information Criterion (WAIC) [59] to identify the psychophysical model that best describes our data (Fig. 3D). This metric evaluates the predictive power of each psychometric model [61] and corrects for the number of parameters to favor parsimony; using this metric, we can compare different models to data and evaluate their goodness of fit. We evaluated three different models: i) Independent - assuming each participant has a single independent JND (and a constant WF), which allows for inter-participant differences in JND posterior distributions ii) Pooled - assuming all participant JNDs arose from a single posterior distribution, and which therefore does not allow for inter-participant differences (this model also features a constant WF), and iii) Variable WF - assuming each participant can have two JNDs. The first JND was calculated using the metabolic data that corresponded to absolute reference costs in the lower half of that participant's reference cost magnitudes, and the second JND was calculated using the metabolic data in the upper half of reference magnitudes. Thus, this model featured a non-constant WF in which the JND varies based on the absolute magnitude of the reference cost. The best model was the independent-JND model with a constant WF (described in *i* above), which obtains the highest WAIC score and is outside the standard error regions of both competing models.


We examined the range of absolute metabolic rates experienced by participants in our protocol and verified that the metabolic rates humans experience while walking with an assistive exoskeleton ($\sim 10\%$ reductions from unassisted walking) were included in this range (Fig. 3E). In conjunction with our constant-WF assumption, this allows the JNDs calculated here to also characterize the perception of energetics when humans are walking with reduced costs due to an exoskeleton.

Discussion

Modern augmentative exoskeletons do not yet provide a metabolic benefit sufficient to exceed the perceptual threshold of human energetics. We demonstrated that the inter-participant average JND of metabolic rate was $22.7\% \pm 17.0\%$. This is substantially greater than the typical reductions obtained using state-of-the-art exoskeletons over the past decade (see Fig. 4) [8]. While some studies have shown metabolic reductions greater than 15% [36, 37, 38], most research has demonstrated more modest reductions. The mean reduction in metabolic rate over the past decade is $\sim 9.6\% \pm 4.5\%$ (averaged from studies in Fig. 4). Based on the inter-participant psychophysical curve obtained in this work, there is a 61% likelihood an average user would perceive a 9.6% change in metabolic rate, when compared to walking without an exoskeleton (50% accuracy would be a random guess). Thus, based on the metabolic rate reductions provided to date [8], these benefits cannot yet be a critical factor in the short-term, conscious perception of exoskeleton use. The perceptual threshold presented in this work (*i.e.* the JND) can act as a useful benchmark for future exoskeletons designed to noticeably improve walking energetic efficiency.

For augmentative exoskeletons to demonstrate value to their wearers, the benefits provided should be perceivable in the short term. Given that state-of-the-art exoskeletons cannot yet exceed the perceptual threshold of metabolic rate (*i.e.* $> 23\%$), this reduction in metabolic rate is not likely to be the driving factor for why users choose to wear these technologies. Given the short-term nature of the trials in this study, it is possible the reduction in metabolic rate is more perceivable over an extended period of use. While this could positively impact user experience, perception over a longer duration may also lead to challenges in experience and adoption. Prior work in economics has demonstrated that a benefit provided in the future is less valuable when compared to a more immediate benefit (*i.e.* temporal discounting) [62, 63, 64, 65, 66, 67, 68]. Thus, we believe exoskeletons will be most successful if the metrics used to develop these technologies are aligned with what is perceivable and valuable to the user in the short term. Understanding if and how longer-term energetic reductions are perceivable, in addition to the impact of temporal discounting, are important avenues of future study.

Metabolic rate reduction is currently the “gold standard” for augmentative exoskeletons, which is supported by its role in the reduction of joint mechanical power, previous biomechanical studies, and its objective measurability. However, our results demonstrate that the current reductions in metabolic rate are not



yet broadly perceptible in the short term. The difficulty of perceiving changes to metabolic rate motivates the consideration of alternative metrics which may be more clearly perceivable by users, including reduction of muscle fatigue [69, 70, 71], peak joint forces in arthritic joints [70, 72, 73], and user preference [74, 75, 76, 77]. The development of perceivable and meaningful metrics to quantify success in future exoskeletons is an important challenge for the field.

Previous work investigating the perception of exertion has shown lower thresholds for exertion during exercise cycling [78]. Haile *et al.* applied the method of adjustment [22] to cycling intensity, arriving at a threshold of 0.15 L/min $\dot{V}O_2$, but did not provide a resting metabolic rate for their participants. Using the resting rate estimates provided in [79], this equates to a JND of $\sim 10\%$. There are several possible explanations for the differences from our results. For example, the method of adjustment can lead to lower JND estimates [80], and is known to be less reliable [81, 82, 83] than forced-choice experiments. Additionally, the exertion levels tested were substantially greater than what was tested in our experiment, and thus might have occurred in the perceptual regime where the Weber Fraction was non-constant. Lastly, their method of moderating exertion used only a single variable (cycling resistance), which is susceptible to confounding factors. That is, cycling resistance will vary proportionally with muscle loading, which can be more easily

sensed through the Golgi tendon organs, mechanoreceptors, and other mechanisms. Thus, any perceptual thresholds calculated using these sensations could be confounded to underestimate the true JND of exertion because the participants could intuit a mapping between the easier-to-perceive cycling resistance and the harder-to-perceive metabolic effort.

Researchers have established that humans will seek energetically optimal gaits, even when metabolic rate changes are far below our estimate of the perceptual threshold for metabolic rate (*i.e.* $\sim 5\%$ rather than 22.7%) [15, 84]. One potential explanation for these differences is that our experiment measures *conscious* perception of changes in metabolic rate, whereas this prior work has allowed for potential subconscious sensing contributions from sensorimotor system and autonomic nervous system [85, 86]. The literature suggests that humans rely on a combination of different afferent signals, such as heart rate or muscular strain, to generate a gestalt perception of exertion in ways that are not yet fully understood [87, 88, 89, 90, 42]. It is also not yet known whether the observed changes in locomotor mechanics that are correlated with metabolic rate are causally linked to those changes.

Participants varied greatly in their ability to perceive changes to their metabolic rate. In this study we investigated whether the JND was more appropriately modeled as a constant value or a person-specific

value. Using the WAIC metric—a modern Bayesian tool for comparing the quality of models—we found that the data were better fit by the model where each participant had their own independent JND (see Fig. 3.D). While the inter-participant mean JND value was greater than the metabolic benefits provided by modern exoskeletons, our participant pool included three participants who had JNDs below 9.6% and thus would likely perceive benefits from these technologies [8] (see Fig. 3.C). Future work is needed to study both the physiological mechanisms that underlie this keener perception of energetics, as well as discovering methods to identify those users who may have better perception.

Our results confirm that the Weber Fraction (WF)—the ratio of the JND (in absolute units of W/kg) to its corresponding reference—is constant with respect to reference magnitude. To this end, we again used the WAIC metric to compare a model with and without a WF dependence on reference magnitude, and found the constant WF model superior. This result indicates that metabolic rate perception in the inter-participant range from ~ 1.5 to 6.6 W/kg (see Fig. 3E) is not near the perceptual extremes where the WF is known to change drastically [50]. The magnitude of this range, in conjunction with the flatness of the WF over this range, also indicates that the perception of metabolic rate penalties is similar to that of metabolic rate improvements.

Limitations

The posterior distributions for those participants with low and high JNDs were differently shaped, reflecting a limit on the maximum metabolic rate changes possible via exoskeleton assistance. The exoskeleton used in this experiment was capable of providing a peak torque of approximately 30 Nm (~ 10 J per stride), which limited the available metabolic rates that could be experienced. The ability to induce a wide array of metabolic rates is important for sampling the psychometric function. To obtain estimates of these functions that have low uncertainty, they must be sampled across both the constant and transitory regions of the psychometric curve [22]. The quality of the measurements is reflected in the posterior distributions for the JND estimates, with high quality measurements resulting in narrow posterior distributions. For participants with smaller JNDs, the limitation on available metabolic rates enabled the sampling of the majority of the relevant areas of the psychometric curve. This allowed us to exclude both excessively large and small estimates for those participants. In contrast, for participants with high JNDs, the imposed energetics spanned a comparatively narrower region of the psychometric function, which only excluded *lower* JNDs.

The posterior distributions for participants with high JNDs was asymmetric, and thus contained greater uncertainty in the upper bound of the threshold. Consequently, any error would likely bias the true JND to be greater than what was measured in this study.

The indirect nature of manipulating energetics via an exoskeleton increased variability in each participant's JND distribution. In conventional psychophysical studies, researchers have more deterministic control over the applied stimulus under investigation. While exoskeletons are known to influence energetics indirectly through several controllable [43, 11, 12, 91] aspects of the torque profile, metabolic rate also depends on many uncontrollable factors that appear noise-like [79, 92, 48]. This added noise results in sub-optimal sampling of the psychophysical curve that reduces certainty in the corresponding JND estimates [49]. This uncertainty is reflected in the width of the posterior distributions of each participant.

The uncertainty of our results also stems from an experimental limitation in how many trials are feasible. Conventional best practice in the psychophysics literature would recommend ~ 300 trials [52] when estimating the underlying psychophysical curve; however, in this study we were able to obtain ~ 100 trials for each participant. The relatively low number of trials was due to the time necessary to obtain responses. In this protocol, participants experienced different metabolic rates in sequence, each of which requires two minutes to estimate the participant's metabolic rate. To obtain the necessary data for this experiment, participants walked during three sessions spread across three days, with each session lasting four hours. This is in contrast to many studies of human perception, which can obtain experimental data without the time delay of the human cardiopulmonary system ($\tau_r = 42$ s [93]). Consequently, the uncertainty of our estimates was increased by approximately 60% [49] due to the lower number of samples, which is reflected in the inter-participant distribution of the JNDs.

We found that despite the uncertainty in JND estimates, these estimates were relatively insensitive to assumptions in our approach. We used a uniform prior distribution in our analysis that encompassed available JNDs between 0% and 70%. We investigated the sensitivity of our results to the bounds of this prior distribution (*i.e.* 0% and 70%). We chose our lower bound to reflect perfect human perception, while the upper bound was informed by the reasonable assumption that a human could consistently detect changes in energetics just under those that result from switching from walking to running (a $\sim 100\%$ change [94]). When the bounds of our uniform prior distribution were changed to [0%, 60%] and [0%, 80%], we found that the

average inter-participant JND estimate shifted from 22.7% to 21.9% and 23.2%, respectively. These small shifts in the mean JND estimate indicate that our approach is robust to the exact shape of our prior distributions, and are thus well-informed by our sampled data.

Participants responded to questions about exertion, but we are unable to know what specifically drove their answers. Our study relies on participants honestly reporting perceived exertion and not confounding this report with other perceptions, which could include perceptions of assistive torque and assistance timing, as well as higher-level perceptions of the helpfulness of the actuation profile. Our study was designed to mitigate these confounding factors. Participants read a predefined script to help elucidate the concepts of metabolic rate and exertion. The prompt was designed using vocabulary consistent with the Borg Scale, used to assess exertion [40, 42, 89]. Additionally, the torque profile was designed to be intentionally complex (see Methods). That is, the participant's metabolic rate was induced by the complex interaction of three controller parameters, obscuring any foreseeable relationship with metabolic rate (*i.e.* "it feels more powerful when it is stronger, which must lower my exertion"). However, if participant's JNDs were affected by additional informative sources, this would also bias the true JND of metabolic perception to be greater than what was estimated.

Conclusion

Motivated by the need to develop augmentative exoskeletons that can realize their potential to impact society, we quantified the human ability to perceive the metabolic impact of these technologies. Participants were able to perceive a 22.7% ($\pm 17.0\%$) change in their metabolic rate with 75% accuracy. Thus, the average user cannot consciously perceive the metabolic benefits from any modern exoskeletons, which may hinder translation and adoption of these technologies. Our results provide a new benchmark for augmentative exoskeletons that will enable perceivable value to their users. The relatively insensitive perception of metabolic rate also suggests that alternative metrics for exoskeleton success, such as reduced muscle fatigue, loading, or user preference, may be more significant to user experience and exoskeleton success.

Availability of data and materials

The datasets generated and/or analysed during the current study are available in the CodeOcean repository, <https://codeocean.com/capsule/8525772/tree> [95].

Competing interests

The authors declare that they have no competing interests.

Author's contributions

RLM, GCT, and EJР designed the study; RLM performed experiment and analyzed data with input from GCT and EJР.; RLM, GCT, and EJР wrote the paper. All authors approved the final version of the manuscript.

Acknowledgements

Text for this section ...

Author details

¹Department of Mechanical Engineering, University of Michigan, 48109 Ann Arbor, USA. ²Department of Electrical Engineering and Computer Science, University of Michigan, 48109 Ann Arbor, USA. ³Robotics Institute, University of Michigan, 48109 Ann Arbor, USA.

References

- Panizzolo, F.A., Galiana, I., Asbeck, A.T., Siviу, C., Schmidt, K., Holt, K.G., Walsh, C.J.: A biologically-inspired multi-joint soft exosuit that can reduce the energy cost of loaded walking. *Journal of NeuroEngineering and Rehabilitation* **13**(1) (2016). doi:[10.1186/s12984-016-0150-9](https://doi.org/10.1186/s12984-016-0150-9)
- Panizzolo, F.A., Freisinger, G.M., Karavas, N., Eckert-Erdheim, A.M., Siviу, C., Long, A., Zifchock, R.A., LaFiandra, M.E., Walsh, C.J.: Metabolic cost adaptations during training with a soft exosuit assisting the hip joint. *Scientific Reports* **9**(1), 1–10 (2019). doi:[10.1038/s41598-019-45914-5](https://doi.org/10.1038/s41598-019-45914-5)
- Kim, J., Lee, G., Heimgartner, R., Revi, D.A., Karavas, N., Nathanson, D., Galiana, I., Eckert-Erdheim, A., Murphy, P., Perry, D., Menard, N., Choe, D.K., Malcolm, P., Walsh, C.J.: Reducing the metabolic rate of walking and running with a versatile, portable exosuit. *Science* **365**(6454), 668–672 (2019). doi:[10.1126/science.aav7536](https://doi.org/10.1126/science.aav7536)
- Mooney, L.M., Lai, C.H., Rouse, E.J.: Design and Characterization of a Biologically Inspired Quasi- Passive Prosthetic Ankle-Foot. 36th Annual International Conference of the IEEE Engineering in Medicine and Biology Society **02139**, 1611–1617 (2014)
- Sawicki, G.S., Ferris, D.P.: Mechanics and energetics of level walking with powered ankle exoskeletons. *Journal of Experimental Biology* **211**(9), 1402–1413 (2008). doi:[10.1242/jeb.009241](https://doi.org/10.1242/jeb.009241)
- Bryan, G.M., Franks, P.W., Klein, S.C., Peuchen, R.J., Collins, S.H.: A hip-knee-ankle exoskeleton emulator for studying gait assistance. *International Journal of Robotics Research* (2020). doi:[10.1177/0278364920961452](https://doi.org/10.1177/0278364920961452)
- Zhu, H., Nesler, C., Divekar, N., Peddinti, V., Gregg, R.: Design principles for compact, backdrivable actuation in partial-assist powered knee orthoses. *IEEE/ASME Transactions on Mechatronics* (2021). early access
- Sawicki, G.S., Beck, O.N., Kang, I., Young, A.J.: The exoskeleton expansion: Improving walking and running economy. *Journal of NeuroEngineering and Rehabilitation* **17**(1), 1–9 (2020). doi:[10.1186/s12984-020-00663-9](https://doi.org/10.1186/s12984-020-00663-9)
- Mooney, L.M., Rouse, E.J., Herr, H.M.: Autonomous exoskeleton reduces metabolic cost of walking. 2014 36th Annual International Conference of the IEEE Engineering in Medicine and Biology Society, EMBC 2014 **11**(1), 3065–3068 (2014). doi:[10.1109/EMBC.2014.6944270.759764](https://doi.org/10.1109/EMBC.2014.6944270.759764)
- Mooney, L.M., Rouse, E.J., Herr, H.M.: Autonomous exoskeleton reduces metabolic cost of human walking during load carriage. *Journal of NeuroEngineering and Rehabilitation* **11**(1), 1–5 (2014). doi:[10.1186/1743-0003-11-151](https://doi.org/10.1186/1743-0003-11-151)
- Malcolm, P., Derave, W., Galle, S., De Clercq, D.: A Simple Exoskeleton That Assists Plantarflexion Can Reduce the Metabolic Cost of Human Walking. *PLoS ONE* **8**(2), 1–7 (2013). doi:[10.1371/journal.pone.0056137](https://doi.org/10.1371/journal.pone.0056137)
- Jackson, R.W., Collins, S.H.: An experimental comparison of the relative benefits of work and torque assistance in ankle exoskeletons. *Journal of Applied Physiology* **119**(5), 541–557 (2015). doi:[10.1152/jappphysiol.01133.2014](https://doi.org/10.1152/jappphysiol.01133.2014)
- Zhang, J., Fiers, P., Witte, K.A., Jackson, R.W., Poggensee, K.L., Atkeson, C.G., Collins, S.H.: Human-in-the-loop optimization of exoskeleton assistance during walking. *Science* **1284**(June), 1280–1284 (2017). doi:[10.1016/j.gaitpost.2011.08.025](https://doi.org/10.1016/j.gaitpost.2011.08.025). The

14. Donelan, J.M., Kram, R., Kuo, A.D.: Mechanical and metabolic determinants of the preferred step width in human walking. *Proceedings of the Royal Society B: Biological Sciences* **268**(1480), 1985–1992 (2001). doi:[10.1098/rspb.2001.1761](https://doi.org/10.1098/rspb.2001.1761)
15. Selinger, J.C., O'Connor, S.M., Wong, J.D., Donelan, J.M.: Humans Can Continuously Optimize Energetic Cost during Walking. *Current Biology* **25**(18), 2452–2456 (2015). doi:[10.1016/j.cub.2015.08.016](https://doi.org/10.1016/j.cub.2015.08.016)
16. Wong, J.D., Selinger, J.C., Donelan, J.M.: Is natural variability in gait sufficient to initiate spontaneous energy optimization in human walking? *Journal of neurophysiology* **121**(5), 1848–1855 (2019). doi:[10.1152/jn.00417.2018](https://doi.org/10.1152/jn.00417.2018)
17. Davis, F.D.: Perceived usefulness, perceived ease of use, and user acceptance of information technology. *MIS Quarterly: Management Information Systems* **13**(3), 319–339 (1989). doi:[10.2307/249008](https://doi.org/10.2307/249008)
18. King, W.R., He, J.: A meta-analysis of the technology acceptance model. *Information and Management* **43**(6), 740–755 (2006). doi:[10.1016/j.im.2006.05.003](https://doi.org/10.1016/j.im.2006.05.003)
19. Oh, S., Ahn, J., Kim, B.: Adoption of broadband internet in korea: the role of experience in building attitudes. *Journal of Information Technology* **18**(4), 267–280 (2003)
20. Hu, P.H., Chau, P., Sheng, O.R., Tam, K.: Examining the technology acceptance model using physician acceptance of telemedicine technology. *J. Manag. Inf. Syst.* **16**, 91–112 (1999)
21. Chuah, S.H.W., Rauschnabel, P.A., Krey, N., Nguyen, B., Ramayah, T., Lade, S.: Wearable technologies: The role of usefulness and visibility in smartwatch adoption. *Computers in Human Behavior* **65**, 276–284 (2016). doi:[10.1016/j.chb.2016.07.047](https://doi.org/10.1016/j.chb.2016.07.047)
22. Kingdom, F., Prins, N.: *Psychophysics: A Practical Introduction* vol. 53, pp. 1689–1699 (2013). doi:[10.1017/CBO9781107415324.004](https://doi.org/10.1017/CBO9781107415324.004). [arXiv:1011.1669v3](https://arxiv.org/abs/1011.1669v3)
23. Gescheider, G.A.: *Psychophysics: the Fundamentals*. Psychology Press, ??? (2013)
24. Wichmann, F.A., Hill, N.J.: The psychometric function: II. Bootstrap-based confidence intervals and sampling. *Perception and Psychophysics* **63**(8), 1314–1329 (2001). doi:[10.3758/BF03194545](https://doi.org/10.3758/BF03194545)
25. Torgerson, W.S.: Theory and methods of scaling. (1958)
26. Stevens, S.S., Galanter, E.H.: Ratio scales and category scales for a dozen perceptual continua. *Journal of experimental psychology* **54**(6), 377 (1957)
27. Stevens, S.S.: On the psychophysical law. *Psychological Review* **64**(3), 153–181 (1957). doi:[10.1037/h0046162](https://doi.org/10.1037/h0046162)
28. BEEBE-CENTER, J.G., WADDELL, D.: A general psychological scale of taste. *The Journal of psychology* **26**, 517–524 (1948). doi:[10.1080/00223980.1948.9917423](https://doi.org/10.1080/00223980.1948.9917423)
29. Brodie, E.E., Ross, H.E.: Sensorimotor mechanisms in weight discrimination. *Perception —& Psychophysics* **36**(5), 477–481 (1984). doi:[10.3758/BF03207502](https://doi.org/10.3758/BF03207502)
30. Shepherd, M.K., Azocar, A.F., Major, M.J., Rouse, E.J.: Amputee perception of prosthetic ankle stiffness during locomotion. *Journal of NeuroEngineering and Rehabilitation* **15**(1), 1–10 (2018). doi:[10.1186/s12984-018-0432-5](https://doi.org/10.1186/s12984-018-0432-5)
31. Shepherd, M.K., Rouse, E.J.: Comparing preference of ankle–foot stiffness in below-knee amputees and prosthetists. *Scientific Reports* **10**(1), 16067 (2020). doi:[10.1038/s41598-020-72131-2](https://doi.org/10.1038/s41598-020-72131-2)
32. Azocar, A.F., Rouse, E.J.: Stiffness Perception During Active Ankle and Knee Movement. *IEEE Transactions on Biomedical Engineering* **64**(12), 2949–2956 (2017)
33. Azocar, A.F., Mooney, L.M., Hargrove, L.J., Rouse, E.J.: Design and Characterization of an Open-source Robotic Leg Prosthesis. *International Conference on Biomedical Robotics and Biomechatronics* (2018)
34. Ackerley, R., Wasling, H.B., Ortiz-Catalan, M., Brånemark, R., Wessberg, J.: Case studies in neuroscience: Sensations elicited and discrimination ability from nerve cuff stimulation in an amputee over time. *Journal of Neurophysiology* **120**(1), 291–295 (2018). doi:[10.1152/jn.00909.2017](https://doi.org/10.1152/jn.00909.2017)
35. Clemente, F., Hakansson, B., Cipriani, C., Wessberg, J., Kulbacka-Ortiz, K., Brånemark, R., Fredén Jansson, K.J., Ortiz-Catalan, M.: Touch and Hearing Mediate Osseoperception. *Scientific Reports* **7**, 1–11 (2017). doi:[10.1038/srep45363](https://doi.org/10.1038/srep45363)
36. Lee, S., Kim, J., Baker, L., Long, A., Karavas, N., Menard, N., Galiana, I., Walsh, C.J.: Autonomous multi-joint soft exosuit with augmentation-power-based control parameter tuning reduces energy cost of loaded walking. *Journal of neuroengineering and rehabilitation* **15**(1), 1–9 (2018)
37. Ding, Y., Kim, M., Kuindersma, S., Walsh, C.J.: Human-in-the-loop optimization of hip assistance with a soft exosuit during walking. *Science Robotics* **3**(15), 1–9 (2018). doi:[10.1126/scirobotics.aar5438](https://doi.org/10.1126/scirobotics.aar5438)
38. Lim, B., Lee, J., Jang, J., Kim, K., Park, Y.J., Seo, K., Shim, Y.: Delayed output feedback control for gait assistance with a robotic hip exoskeleton. *IEEE Transactions on Robotics* **35**(4), 1055–1062 (2019)
39. Collins, S.H., Zhang, J., Poggensee, K.L., Witte, K.A., Jackson, R.W., Fiers, P., Atkeson, C.G.: Supplementary Materials for Human-in-the-loop optimization of exoskeleton assistance during walking. *Science* **356**(6344), 1280–1284 (2017). doi:[10.1126/science.aal5054](https://doi.org/10.1126/science.aal5054)
40. Borg, E., Kaijser, L.: A comparison between three rating scales for perceived exertion and two different work tests. *Scandinavian Journal of Medicine and Science in Sports* **16**(1), 57–69 (2006). doi:[10.1111/j.1600-0838.2005.00448.x](https://doi.org/10.1111/j.1600-0838.2005.00448.x)
41. Borg, E.: So What's that On a Scale of 1 to 10. *Proceedings of the 24th International Congress of Vexillology (August 2011)*, 988–995 (2011)
42. Borg, E.: On Perceived Exertion and Its Measurement, (2007). <http://su.diva-portal.org/smash/get/diva2:197216/FULLTEXT01>
43. Koller, J.R., Gates, D.H., Ferris, D.P., Remy, C.D.: Confidence in the curve: Establishing instantaneous cost mapping techniques using bilateral ankle exoskeletons. *Journal of Applied Physiology* **122**(2), 242–252 (2016). doi:[10.1152/jappphysiol.00710.2016](https://doi.org/10.1152/jappphysiol.00710.2016)
44. Samuel, G., Philippe, M., Dirk, D.C.: 2D Parameter sweep of bilateral exoskeleton actuation, pp. 2–3. *Dynamic Walking, ???* (2014)
45. Kang, I., Molinaro, D., Duggal, S., Chen, Y., Kunapuli, P., Young, A.: Real-time gait phase estimation for robotic hip exoskeleton control during multimodal locomotion. *IEEE Robotics and Automation Letters* **6**(2), 3491–3497 (2021). doi:[10.1109/LRA.2021.3062562](https://doi.org/10.1109/LRA.2021.3062562)
46. Camargo, J., Flanagan, W., Csomay-Shanklin, N., Kanwar, B., Young, A.: A Machine Learning Strategy for Locomotion Classification and Parameter Estimation Using Fusion of Wearable Sensors. *IEEE Transactions on Biomedical Engineering* **68**(5), 1569–1578 (2021). doi:[10.1109/TBME.2021.3065809](https://doi.org/10.1109/TBME.2021.3065809)
47. Selinger, J.C., Donelan, J.M.: Estimating instantaneous energetic cost during non-steady-state gait. *Journal of Applied Physiology* **117**(11), 1406–1415 (2014). doi:[10.1152/jappphysiol.00445.2014](https://doi.org/10.1152/jappphysiol.00445.2014)
48. Guidetti, L., Bolletta, F., Gallotta, M.C., Baldari, C., Meucci, M., Emerenziani, G.P.: Validity, reliability and minimum detectable change of COSMED K5 portable gas exchange system in breath-by-breath mode. *PLoS ONE* **13**(12), 1–12 (2018). doi:[10.1371/journal.pone.0209925](https://doi.org/10.1371/journal.pone.0209925)
49. Medrano, R.L., Thomas, G.C., Rouse, E.: Methods for Measuring the Just Noticeable Difference for Variable Stimuli: Implications for Perception of Metabolic Rate with Exoskeleton Assistance. *Proceedings of the IEEE RAS and EMBS International Conference on Biomedical Robotics and Biomechatronics 2020-November*, 483–490 (2020). doi:[10.1109/BioRob49111.2020.9224374](https://doi.org/10.1109/BioRob49111.2020.9224374)
50. Norwich, K.H.: On the theory of Weber fractions. *Perception & Psychophysics* **42**(3), 286–298 (1987). doi:[10.3758/BF03203081](https://doi.org/10.3758/BF03203081)
51. Gescheider, G.a.: *Psychophysical Measurement of Thresholds* (1985)
52. García-Pérez, M.A., Alcalá-Quintana, R.: Sampling plans for fitting the psychometric function. *Spanish Journal of Psychology* **8**(2), 256–289 (2005). doi:[10.1017/S113874160000514X](https://doi.org/10.1017/S113874160000514X)
53. Kuroda, T., Hasuo, E.: The very first step to start psychophysical experiments. *Acoustical Science and Technology* **35**(1), 1–9 (2013). doi:[10.1250/ast.35.1](https://doi.org/10.1250/ast.35.1)
54. Kuss, M., Jäkel, F., Wichmann, F.A.: Bayesian inference for psychometric functions. *Journal of Vision* **5**(5), 478–492 (2005). doi:[10.1167/5.5.8](https://doi.org/10.1167/5.5.8)
55. Wichmann, F.A., Jäkel, F.: *Methods in Psychophysics*, pp. 1–42. Wiley, ??? (2018). doi:[10.1002/9781119170174.epcn507](https://doi.org/10.1002/9781119170174.epcn507)
56. Prins, N., Kingdom, F.A.A.: Applying the Model-Comparison Approach to Test Specific Research Hypotheses in Psychophysical Research Using the Palamedes Toolbox. *Frontiers in Psychology* **9**, 1250 (2018). doi:[10.3389/fpsyg.2018.01250](https://doi.org/10.3389/fpsyg.2018.01250)

57. Salvatier, J., Wiecki, T.V., Fonnesebeck, C.: Probabilistic programming in python using pymc3. *PeerJ Computer Science* **2**, 55 (2016)
58. Hoffman, M.D., Gelman, A.: The no-U-turn sampler: Adaptively setting path lengths in Hamiltonian Monte Carlo. *Journal of Machine Learning Research* **15**(2008), 1593–1623 (2014). [1111.4246](https://doi.org/10.1111.4246)
59. Vehtari, A., Gelman, A., Gabry, J.: Practical Bayesian model evaluation using leave-one-out cross-validation and WAIC. *Statistics and Computing* **27**(5), 1413–1432 (2017). doi:[10.1007/s11222-016-9696-4](https://doi.org/10.1007/s11222-016-9696-4). [1507.04544](https://doi.org/10.1507.04544)
60. Das Gupta, S., Bobbert, M.F., Kistemaker, D.A.: The Metabolic Cost of Walking in healthy young and older adults – A Systematic Review and Meta Analysis. *Scientific Reports* **9**(1), 1–10 (2019). doi:[10.1038/s41598-019-45602-4](https://doi.org/10.1038/s41598-019-45602-4)
61. Gelman, A., Hwang, J., Vehtari, A.: Understanding predictive information criteria for Bayesian models. *Statistics and Computing* **24**(6), 997–1016 (2014). doi:[10.1007/s11222-013-9416-2](https://doi.org/10.1007/s11222-013-9416-2). [1307.5928](https://doi.org/10.1307.5928)
62. Zandstra, E.H., Miyapuram, K.P., Tobler, P.N.: Understanding Consumer Decisions Using Behavioral Economics vol. 202, 1st edn., pp. 197–211. Elsevier B.V., ??? (2013). doi:[10.1016/B978-0-444-62604-2.00012-5](https://doi.org/10.1016/B978-0-444-62604-2.00012-5). <http://dx.doi.org/10.1016/B978-0-444-62604-2.00012-5>
63. Green, L., Fristoe, N., Myerson, J.: Temporal discounting and preference reversals in choice between delayed outcomes. *Psychonomic Bulletin & Review* **1**(3), 383–389 (1994). doi:[10.3758/BF03213979](https://doi.org/10.3758/BF03213979)
64. Green, L., Myerson, J., McFadden, E.: Rate of temporal discounting decreases with amount of reward. *Memory and Cognition* **25**(5), 715–723 (1997). doi:[10.3758/BF03211314](https://doi.org/10.3758/BF03211314)
65. Simpson, C.A., Vuchinich, R.E.: Reliability of a measure of temporal discounting. *Psychological Record* **50**(1), 3–16 (2000). doi:[10.1007/BF03395339](https://doi.org/10.1007/BF03395339)
66. Critchfield, T.S., Kollins, S.H.: Temporal Discounting: Basic Research and the Analysis of Socially Important Behavior. *Journal of Applied Behavior Analysis* **34**(1), 101–122 (2001). doi:[10.1901/jaba.2001.34-101](https://doi.org/10.1901/jaba.2001.34-101)
67. Schultz, W.: Subjective neuronal coding of reward: Temporal value discounting and risk. *European Journal of Neuroscience* **31**(12), 2124–2135 (2010). doi:[10.1111/j.1460-9568.2010.07282.x](https://doi.org/10.1111/j.1460-9568.2010.07282.x)
68. Bos, W.V.D., McClure, S.M.: Towards a general model of temporal discounting. *Journal of the Experimental Analysis of Behavior* **99**(1), 58–73 (2013). doi:[10.1002/jeab.6](https://doi.org/10.1002/jeab.6)
69. McDonald, K.: Multi-objective prioritization in human walking (2018)
70. Ackermann, M., Van Den Bogert, A.J.: Optimality Principles for Model-Based Prediction of Human Gait. *Physical Therapy* **31**(January 2008), 1–8 (2011). doi:[10.1016/j.jbiomech.2009.12.012](https://doi.org/10.1016/j.jbiomech.2009.12.012). [Optimality](https://doi.org/10.1016/j.jbiomech.2009.12.012)
71. Crowell, H.P., Kanagaki, G.B., O'donovan, M.P., Haynes, C.A., Park, J.-H., Neugebauer, J.M., Hennessy, E.R., Boynton, A.C., Mitchell, B., Tweedell, A.J., Girolamo, H.J.: Methodologies for Evaluating the Effects of Physical Augmentation Technologies on Soldier Performance. US Army Research Laboratory Aberdeen Proving Ground United States (May) (2018). doi:[10.13140/RG.2.2.13662.48961](https://doi.org/10.13140/RG.2.2.13662.48961)
72. Hunter, L.C., Hendrix, E.C., Dean, J.C.: The cost of walking downhill: Is the preferred gait energetically optimal? *Journal of Biomechanics* **43**(10), 1910–1915 (2010). doi:[10.1016/j.jbiomech.2010.03.030](https://doi.org/10.1016/j.jbiomech.2010.03.030)
73. Medrano, R.L., Rouse, E.J., Thomas, G.C.: Biological Joint Loading and Exoskeleton Design. *IEEE Transactions on Medical Robotics and Bionics*, 1–4
74. Thatte, N., Duan, H., Geyer, H.: A sample-efficient black-box optimizer to train policies for human-in-the-loop systems with user preferences. *IEEE Robotics and Automation Letters* **2**(2), 993–1000 (2017)
75. Ingraham, K.A., Remy, C.D., Rouse, E.J.: User preference of applied torque characteristics for bilateral powered ankle exoskeletons. *Proceedings of the IEEE RAS and EMBS International Conference on Biomedical Robotics and Biomechatronics 2020-November*, 839–845 (2020). doi:[10.1109/BioRob49111.2020.9224358](https://doi.org/10.1109/BioRob49111.2020.9224358)
76. Shepherd, M.K., Azocar, A.F., Major, M.J., Rouse, E.J.: Amputee perception of prosthetic ankle stiffness during locomotion. *Journal of NeuroEngineering and Rehabilitation* **15**(1), 1–10 (2018). doi:[10.1186/s12984-018-0432-5](https://doi.org/10.1186/s12984-018-0432-5)
77. Tucker, M., Novoseller, E., Kann, C., Sui, Y., Yue, Y., Burdick, J., Ames, A.D.: Preference-Based Learning for Exoskeleton Gait Optimization. 2020 IEEE International Conference on Robotics and Automation (ICRA) (2019). [1909.12316](https://doi.org/10.1109/ICRA.2019.8792032)
78. Haile, L., Robertson, R.J., Nagle, E.F., Krause, M.P., Gallagher, M., Ledezma, C.M., Wisniewski, K.S., Shafer, A.B., Goss, F.L.: Just noticeable difference in perception of physical exertion during cycle exercise in young adult men and women. *European Journal of Applied Physiology* **113**(4), 877–885 (2013). doi:[10.1007/s00421-012-2497-3](https://doi.org/10.1007/s00421-012-2497-3)
79. Davidson, A., Gardinier, E.S., Gates, D.H.: Within and between-day reliability of energetic cost measures during treadmill walking. *Cogent Engineering* **3**(1), 1–7 (2016). doi:[10.1080/23311916.2016.1251028](https://doi.org/10.1080/23311916.2016.1251028)
80. Wier, C.C., Jesteadt, W., Green, D.M.: A comparison of method-of-adjustment and forced-choice procedures in frequency discrimination. *Perception & Psychophysics* **19**(1), 75–79 (1976). doi:[10.3758/BF03199389](https://doi.org/10.3758/BF03199389)
81. Stevens, S.S.: PROBLEMS AND METHODS OF PSYCHOPHYSICS 1. *Psychological Bulletin* **55**(4) (1958)
82. Cardozo, B.L.: Adjusting the Method of Adjustment: SD vs DL. *The Journal of the Acoustical Society of America* **37**(5), 786–792 (1965). doi:[10.1121/1.1909439](https://doi.org/10.1121/1.1909439)
83. Pelli, D.G., Farell, B.: Chapter 29 - Psychophysical Methods. *Handbook of Optics*, 1–13 (1996)
84. Abram, S.J., Selinger, J.C., Donelan, J.M.: Energy optimization is a major objective in the real-time control of step width in human walking. *Journal of Biomechanics* **91**, 85–91 (2019). doi:[10.1016/j.jbiomech.2019.05.010](https://doi.org/10.1016/j.jbiomech.2019.05.010)
85. Wong, J.D., O'Connor, S.M., Selinger, J.C., Donelan, J.M.: Contribution of blood oxygen and carbon dioxide sensing to the energetic optimization of human walking. *Journal of Neurophysiology*, 00195–2017 (2017). doi:[10.1152/jn.00195.2017](https://doi.org/10.1152/jn.00195.2017)
86. Selinger, J.C., Wong, J.D., Simha, S.N., Donelan, J.M.: How people initiate energy optimization and converge on their optimal gaits. *The Journal of Experimental Biology* (September), 198234 (2019). doi:[10.1242/jeb.198234](https://doi.org/10.1242/jeb.198234)
87. Hampson, D.B., Clair Gibson, A.S., Lambert, M.I., Noakes, T.D.: The influence of sensory cues on the perception of exertion during exercise and central regulation of exercise performance. *Sports Medicine* **31**(13), 935–952 (2001). doi:[10.2165/00007256.200131130-00004](https://doi.org/10.2165/00007256.200131130-00004)
88. Gibson, A.S.C., Baden, D.A., Lambert, M.I., Lambert, E.V., Harley, X.R., Hampson, D., Russell, V.A., Noakes, T.D.: The Conscious Perception of the Sensation of Fatigue. *Sports Medicine* **33**(3), 1–10 (2003)
89. Scherr, J., Wolfarth, B., Christle, J.W., Pressler, A., Wagenpfeil, S., Halle, M.: Associations between Borg's rating of perceived exertion and physiological measures of exercise intensity. *European Journal of Applied Physiology* **113**(1), 147–155 (2013). doi:[10.1007/s00421-012-2421-x](https://doi.org/10.1007/s00421-012-2421-x)
90. Borg, G.A.V.: Psychophysical bases of perceived exertion (1982)
91. Galle, S., Malcolm, P., Collins, S.H., De Clercq, D.: Reducing the metabolic cost of walking with an ankle exoskeleton: interaction between actuation timing and power. *Journal of NeuroEngineering and Rehabilitation* **14**(1), 1–16 (2017). doi:[10.1186/s12984-017-0235-0](https://doi.org/10.1186/s12984-017-0235-0)
92. Hangen, H., Melanson, E., Tran, Z., Kearney, J. T., Hill, J.O.: Variability of resting metabolic rate. *American Journal of Clinical Nutrition* **78**, 1141–1144 (2003)
93. Zhang, J., Cheah, C.C., Collins, S.H.: Torque Control in Legged Locomotion, 1st edn., pp. 347–400. Elsevier Inc., ??? (2017). doi:[10.1016/B978-0-12-803766-9.00007-5](https://doi.org/10.1016/B978-0-12-803766-9.00007-5). <http://dx.doi.org/10.1016/B978-0-12-803766-9.00007-5>
94. Teunissen, L.P.J., Grabowski, A., Kram, R.: Effects of independently altering body weight and body mass on the metabolic cost of running. *Journal of Experimental Biology* **210**(24), 4418–4427 (2007). doi:[10.1242/jeb.004481](https://doi.org/10.1242/jeb.004481)
95. Medrano, R.L., Thomas, G.C., Rouse, E.J.: Can Humans Perceive the Metabolic Benefit Provided by Augmentative Exoskeletons? [Source Code]. <https://doi.org/10.24433/CO.8128032.v1>